

Patient Name: \_\_\_\_\_ Date: \_\_\_\_\_ DOB: \_\_\_\_\_

Telephone: \_\_\_\_\_ Address: \_\_\_\_\_

Fax Order 402-408-0020 / **MARK APPROPRIATE COMPOUND**

**Commonly Used for Dermatitis - 30gm**

\_\_\_\_\_ Doxepin HCL 5%/ Flurbiprofen 5%/ Lidocaine 2%/Triamcinolone Acetonide Micronized 0.1% in Lipoderm

\_\_\_\_\_ Sig: Apply 2-3 times daily

**Commonly Used for Scars - 30gm**

\_\_\_\_\_ (Post Surgery)Tamoxifen Citrate 0.1% / in Pracisil

\_\_\_\_\_ (Keloid Scar) Betamethasone 0.5%, Loratadine 1% Topical Gel in Pracasil

\_\_\_\_\_ (Stretch Marks) Topiramate 2.5% / Aloe Vera 0.5%/Pracisil

\_\_\_\_\_ Sig: Apply 2-3 times daily

**Commonly Used for Shingles/Postherpetic Neuralgia -30gm**

\_\_\_\_\_ Deoxy-D-Glucose(2) 0.2%, Gabapentin 6%, Flurbiprofen 10%, Amitriptyline HCL 2% Tetracaine HCL 1% Topical in Lipoderm

\_\_\_\_\_ Sig: Apply 4-5 times daily during active outbreak. Once lesions clear, apply QID until pain decreases then TID until pain is gone.

**Commonly Used for Eczema or Psoriasis - 30gm**

\_\_\_\_\_ Cyanocobalamin 0.7% with Tranilast 0.5%

\_\_\_\_\_ Clobetasol Soln 50cc / camphor0.5% / Menthol0.5%/ Cerave Cream 16oz. QS

\_\_\_\_\_ Desonide 0.075% in Lipoderm

**Commonly Used for Nail Fungus - 10ml**

\_\_\_\_\_ Itraconazole 1%, Flurbiprofen 2%, DMSO

\_\_\_\_\_ Terbinafine 3%, Fluconazole 3%, Flurbiprofen 2% in DMSO

\_\_\_\_\_ Sig: Apply 3-4 times daily as directed

**Commonly Used for Rosacea - 30g**

\_\_\_\_\_ Metronidazole 1% / Niacinamide 4% Gel

\_\_\_\_\_ Ivermectin 1% / Azelaic Acid 15%

**Commonly Used for Warts - 10ml**

\_\_\_\_\_ Salicylic Acid 20%, Fluorouracil 5%, Retinoic Acid 0.1%

\_\_\_\_\_ Salicylic Acid 30%, Retinoic Acid 0.1% (Recommended for children under 8)

\_\_\_\_\_ Glycolic Acid 15%, Salicylic Acid 2% in Topical Gel

\_\_\_\_\_ Cantharidin 0.7%

\_\_\_\_\_ Sig: Apply daily at bedtime cover with occlusive dressing

**Commonly Used for Keratosis Pilaris - 30gm - Use as directed**

\_\_\_\_\_ Salicylic Acid 8%, Glycolic Acid 5% in Amlactin Lotion

**Commonly Used for Skin Brightening - 30gm - Use as directed**

\_\_\_\_\_ Tretinoin 0.1%, Fluocinolone 0.01%, Hydroquinone 5% gel

**Commonly Used for Age Spots - 30gm - use as directed**

\_\_\_\_\_ Hydroquinone 11%/ Kojic Acid 4% Gel

\_\_\_\_\_ Hydroquinone Gel 4% - 6% - 8%

**Commonly Used for Diaper Rash -65gm - Use as directed**

\_\_\_\_\_ Cholestyramine in Aquaphor 12gm / 53gm

**Commonly Used for Precancerous Spots - 15gm - Use as directed**

\_\_\_\_\_ 5-Fluorouracil 2% Glycolic Acid 2.5% Cream

**Commonly Used for Lip Moisturizer 3g - Use as directed**

\_\_\_\_\_ 5-Fluorouracil 2% Lip Balm

**Commonly Used for Acne -30gm - Use as directed**

\_\_\_\_\_ Clindamycin 0.6gm / Sulfacetamide 10% / Sulfur 5%

**Commonly Used for Seborrheic Dermatitis -30gm**

\_\_\_\_\_ Fluocinolone Acetonide 0.01% - Cyanocobalamin 0.7%

\_\_\_\_\_ Sig: Use once daily, as directed

M.D. Signature: \_\_\_\_\_

M.D. Name: \_\_\_\_\_ Date: \_\_\_\_\_